NEW HAMPSHIRE WATER SUPPLY AND POLLUTION CONTROL COMMISSION LAKE TROPHIC DATA

MORPHOMETRIC:		
LAKE Clark Pond	LAKE AREA (HA)	55.04
TOWNCanaan		M) 13.4
COUNTYGrafton		5.1
RIVER BASIN <u>Connecticut</u>	VOLUME (M ³)	2,815,000
LATITUDE 43° 44'N	MUD SURFACE ARE	A (HA)55.08
LONGITUDE 72° 04'W	RELATIVE DEPTH	1.6
ELEVATION (FT) <u>1043</u>	SHORE CONFIGURA	TION1.41
SHORE LENGTH (M) 3700	AREAL WATER LOA	D (M/YR) <u>6.81</u>
WATERSHED AREA (HA)777.0	FLUSHING RATE (YR ⁻¹) 1.3
% WATERSHED PONDED0_4%	PHOSPHORUS RETE	NTION COEFF. <u>0.61</u>
BIOLOGICAL: DATE	14 FEB 1986	30 JUL 1985
DOM. PHYTOPLANKTON (% total) ¹	Asterionella (70%)	Chrysosphaerella (40%)
2	Synura (20%)	Asterionella (20%)
NUMBER OF ALGAL GENERA	7	11
SPECIES DIVERSITY		2.48
CHLOROPHYLL <u>a</u> (μg/L)		5.62
DOM. ZOOPLANKTON (% total) 1	Kellicottia (50%)	No sample
2	Keratella (35%)	
ROTIFERS/LITER	129	
MICROCRUSTACEA/LITER	8	
TOTAL ZOOPLANK. CNTS (cells/L)	142	
VASCULAR PLANT ABUNDANCE		Common
DOMINANT VASCULAR PLANTS 1		Eriocaulon
2		Utricularia
3		Nymphaea
SECCHI DISK TRANSPARENCY (M)		~ 4.5
BOTTOM DISS. OXYGEN (mg/L)	3.0	0.6
SEDIMENT: % ORGANIC MATTER		
LAKE TYPE: A natural pond.		
SUMMER THERMAL STRATIFICATION:	YES X NO WEAK	
IF YES, VOLUME OF HYPOLIMNION	294,000 (m³) TH	ERMOCLINE DEPTH 5.0 (m)

CHEMICAL: (mg/L unless	indicated o	therwise)	LAKE: Clark	Pond	
	WIN.	TER		SUMMER	
DATE	14 FEB	1986	30	JUL 1985	
DEPTH (M)	4.0	8.0	2.0	6.5	11.0
pH (UNITS)	6.5	6.4	7.0	6.4	6.2
ALKALINITY (I. P.)	6.0	6.7	5.6	5.2	6.4
ALKALINITY (F.E.P.)	7.8	8.1	7.3	6.9	8.0
NITRITE+NITRATE NITROGEN			< 0.05		< 0.05
TOTAL KJELDAHL NITROGEN			0.50		0.36
TOTAL PHOSPHORUS	NR	NR	0.004	0.010	0.013
SPEC. CONDUCT. (μMhos/cm)	36.2	37.5	31.5	30.9	33.2
APPARENT COLOR (UNITS)	30	30	20	25	30
TRUE COLOR (440 nm)(UNITS)	NR	NR	18	16	25
MAGNESIUM			0.56		
CALCIUM			2.9		
SODIUM			1		
POTASSIUM			< 0.5		
CHLORIDE			< 2		< 2
TN : TP			125		28
INORG-N : INORG-P					
[Mg+Ca] : [Na+K]					
CALCITE SATURATION INDEX			3.1		


* = NOT DEFENSIBLE

NR = NO RESULT

TROPHIC CLASSIFICATION: 1985	D.O.		PLANT ABUND.	CHL <u>a</u>	TOTAL PTS.	TROPHIC CLASS.
CLASSIFICATION POINTS:	4	1	2	1	8	Meso.

COMMENTS:

- 1. Good boat launch with ample parking, although water was shallow.
- 2. About 6 cottages located along the shoreline.
- Road to pond was not plowed in the winter, access was by crosscountry skis.




FIELD DATA SHEET

WATER BODY	Clark Pond		TOWN	Canaan		ВΥ	WSPCC	
DATE COLLEC	TED <u>30 Jul</u>	y 1985	WE	EATHER SU	unny, warm,		ndy	
STATION	DEPTH (M)	TEMP. (°C)	*DISSOLVED OXYGEN	OXYGEN: % SATURATION				
DEEP SPOT	0.1	23.7	8.2	98%				
	1.0	23.6	8.1	97%				
	2.0	23.6	8.1	97%				
	3.0	23.4	8.1	97%				
	4.0	21.9	8.5	97%				
	5.0	17.6	9.7	102%				
	6.0	14.0	7.4	74%				
	7.0	11.8	6.4	69%				
	8.0	9.5	4.8	42%				
	9.0	8.2	3.2	28%				
	10.0	7.9	2.0	17%				
	11.0	7.6	1.5	13%				
	12.0	7.4	0.8	7%				
	13.0	7.4	0.6	5%				
CECCHI DICK (M) 1. E							

SECCHI	DISK ((M) _	4.5	5
BOTTOM	DEPTH	(M)	13.	.0
TIME			1330	hrs.

COMMENTS: 1. Secchi disk transparency is approximate; Kemmerer bottle was visible at 4 meters (forgot Secchi disk) and water was very choppy from high winds.

^{*} Dissolved oxygen values in mg/L



AQUATIC PLANT SURVEY

Key	GENERIC PLAN	ABUNDANCE	
Р	Pontederia cordata	Pickerelweed	Scattered
С	Cyperaceae	Non-flowering sedge	Scattered
L	Lobelia dortmanna	Water lobelia	Scattered
t	Lysimachia terrestris	Swampcandle	Scattered
E	Eriocaulon septangulare	Pipewort	Common
S	Sparganium	Bur Reed	Scattered
Н	Equisetum	Horsetail	Scattered
D	Dulichium arundinaceum	Three-way sedge	Scattered
W	Potamogeton	Pondweed	Scattered
Υ	Nuphar	Yellow water lily	Scattered
d	Drosera	Sundew	Sparse
b	Scirpus	Bulrush	Scattered
N	Nymphaea	White water lily	Common
В	Brasenia schreberi	Water shield	Scattered
U	Utricularia	Bladderwort	Common
Т	Typha	Cattail	Sparse
е	Eleocharis	Spike rush	Sparse
Х		Sterile thread-like leaves	Scattered
G	Gramineae	Grass family	Scattered
·			

GENERAL OBSERVATIONS:

1.Plants were abundant in the northern coves. The rest of the shoreline had scattered to common plants in the near shore area only - they did not extend out into the lake and were not a nuisance problem.

OVERALL ABUNDANCE Common

2.Two loons were observed.